Universität Koblenz-Landau

FB 4 Informatik

Prof. Dr. Viorica Sofronie-Stokkermans Dipl. Inf. Markus Bender

November 12, 2012

Exercises for "Decision Procedures for Verification" Exercise sheet 5

Exercise 5.1:

Let $\Sigma = (\Omega, \Pi)$ be a signature, where $\Omega = \{f/3, g/1, a/0, b/0\}$ and $\Pi = \{p/2\}$; let X be the set of variables $\{x, y, z\}$. Which of the following expressions are terms over Σ and X, which are atoms/literals/clauses/formulae (in first-order logic with equality), which are neither?

- (a) $\neg p(g(a), f(x, y, g(a)))$
- (b) $f(x, x, x) \approx x$
- (c) $p(f(x, x, a), x) \vee p(a, b)$
- (d) $p(\neg g(x), g(y))$
- (e) $\neg p(f(x, y, y))$
- (f) $\neg p(f(x,y),y) \lor p(x,y)$
- (g) $p(a,b) \wedge p(x,y) \wedge y$
- (h) $\exists y(\neg p(f(y,y,y),y))$
- (i) $\forall x \forall y (f(p(x,y),x,x) \approx g(x))$

Exercise 5.2:

Let $\Sigma = (S, \Omega, \Pi)$ be a many-sorted signature, where $S = \{\text{int}, \text{list}\}$, $\Omega = \{\text{cons}, \text{car}, \text{cdr}, \text{nil}, b\}$ and $\Pi = \{p\}$ with the following arities:

```
a(\mathsf{cons}) = \mathsf{int}, \mathsf{list} \to \mathsf{list} \qquad a(\mathsf{car}) = \mathsf{list} \to \mathsf{int} \quad a(\mathsf{cdr}) = \mathsf{list} \to \mathsf{list}
```

 $a(\mathsf{nil}) = \to \mathsf{list}$ (i.e. nil is a constant of sort list)

 $a(b) = \rightarrow \text{ int }$ (i.e. b is a constant of sort int)

a(p) = int, list.

Let X_{int} be the set of variables of sort int containing $\{i, j, k\}$, and let X_{list} be the set of variables of sort list containing $\{x, y, z\}$. Let $X = \{X_{\text{int}}, X_{\text{list}}\}$. Which of the following expressions are terms over Σ and X, which are atoms/literals/clauses/formulae¹, which are neither?

- (a) cons(cons(b, nil), nil)
- (b) cons(b, cons(b, nil))
- (c) $\neg p(b, \mathsf{cons}(b, \mathsf{cons}(b, \mathsf{nil})))$

¹In first-order logic with equality, where equality between terms of sort int is \approx_i and equality between terms of sort list is \approx_i .

```
(d) \neg p(\mathsf{cons}(b, \mathsf{nil}), \mathsf{cons}(b, \mathsf{cons}(b, \mathsf{nil})))
```

(e)
$$cons(b, cons(b, nil)) \approx_l cons(cons(x, b), nil)$$

- (f) $cons(i, cons(b, nil)) \approx j$
- (g) $p(\neg \mathsf{car}(x), x)$
- (h) $\neg p(\mathsf{car}(x), x) \lor p(j, \mathsf{cons}(j, x))$
- (i) $\neg p(b, x) \lor p(b, cons(b, x)) \lor b$
- (j) $\forall i : \mathsf{int}, \forall x : \mathsf{list} \; (\mathsf{cons}(\mathsf{car}(x), \mathsf{cdr}(x)) \approx_l x)$
- (k) $\exists i : \mathsf{int}, \forall y : \mathsf{list} (\mathsf{cons}(b, p(x, y)) \approx_l \mathsf{cdr}(y))$

Exercise 5.3:

Compute the results of the following substitutions:

```
(a) f(g(x), x)[g(a)/x]
```

- (b) p(f(y,x),g(x))[x/y]
- (c) $\forall y (p(f(y,x),g(y)))[x/y]$
- (d) $\forall y(p(f(y,x),x))[y/x]$
- (e) $\forall y(p(f(z,g(y)),g(x)) \vee \exists z(g(z) \approx y))[g(b)/z]$
- (f) $\exists y (f(x,y) \approx x \rightarrow \forall x (f(x,y) \approx x)) [g(y)/y, g(z)/x]$

Please submit your solution until Monday, November 19, 2012 at 9:00. Joint solutions prepared by up to three persons are allowed. Please do not forget to write your name on your solution. Submission possibilities:

- By e-mail to mbender@uni-koblenz.de with the keyword "Homework DP" in the subject.
- Put it in the box in front of Room B 222.