

Collection of exercises: Part 2

Exercise 1. Check the satisfiability of the following formulae using the DAG version of the Congruence Closure algorithm presented in the class:

1. $f(a, b) \approx f(b, a) \wedge f(c, a) \not\approx f(b, c)$
2. $f(g(a)) \approx g(f(a)) \wedge f(g(f(b))) \approx a \wedge f(b) \approx a \wedge g(f(a)) \not\approx a$
3. $f(f(f(a))) \approx f(a) \wedge f(f(a)) \approx a \wedge f(a) \not\approx a$

Exercise 2.

(1a) Check the satisfiability over \mathbb{Z} of the following set of constraints in positive difference logic using the method presented in the lecture. In case of satisfiability find a satisfiable assignment.

- (a) $x - y \leq 4 \wedge y - z \leq 2 \wedge x - z \leq 2 \wedge z - x \leq -3$
- (b) $x - y \leq 4 \wedge y - z \leq 0 \wedge x - z \leq 4 \wedge z - x \leq -3 \wedge x - u \leq -4$

(1b) Check the satisfiability over \mathbb{Z} of the following set of constraints in difference logic using the method presented in the lecture. In case of satisfiability find a satisfiable assignment.

- (a) $x - y \leq 4 \wedge y - z \leq 0 \wedge x - z < 4 \wedge z - x \leq -3 \wedge x - u \leq -4$
- (a) $x - y \leq 4 \wedge y - z \leq 0 \wedge x - z < 4 \wedge z - x < -3 \wedge x - u \leq -4$

(2a) Check the satisfiability over \mathbb{Q} of the following sets of constraints in positive difference logic. In case of satisfiability find a satisfiable assignment.

- (a) $x - y \leq 5 \wedge y - u \leq 4 \wedge x - z \leq -1 \wedge z - x \leq 1.$
- (b) $x - y \leq 5 \wedge y - u \leq 4 \wedge x - z \leq -1 \wedge z - x \leq 1 \wedge z - y \leq -5.$

(2a) Check the satisfiability over \mathbb{Q} of the following sets of constraints in difference logic. In case of satisfiability find a satisfiable assignment.

- (a) $x - y \leq 5 \wedge y - u \leq 4 \wedge x - z < -1 \wedge z - x \leq 1.$
- (b) $x - y \leq 5 \wedge y - u \leq 4 \wedge x - z < -0.5 \wedge z - x < 1 \wedge z - y \leq -5.$

Exercise 3. Use the Nelson-Oppen procedure for checking the satisfiability of the following formulae:

1. $1 \leq c \wedge c \leq 3 \wedge f(c) \not\approx f(1) \wedge f(c) \not\approx f(3) \wedge f(1) \not\approx f(2)$
in the combination $LI(\mathbb{Z}) \cup UIF_{\{f\}}$.
2. $f(c) \approx c + d \wedge c \leq d + e \wedge c + e \leq d \wedge d = 1 \wedge f(c) \not\approx f(2)$
in the combination $LI(\mathbb{Z}) \cup UIF_{\{f\}}$.
3. $c + d \approx e \wedge f(e) \approx e \wedge f(c + d) \not\approx e$
in the combination $LI(\mathbb{Q}) \cup UIF_{\{f\}}$.

Exercise 4. Let $\mathcal{T} = LI(\mathbb{Q})$, and let $Q := x \geq 1, R := x \leq y, P := x + x \leq 2$. Use a DPLL(\mathcal{T}) method to check the satisfiability w.r.t. \mathcal{T} of the following set of clauses:

$$\begin{array}{ll} (C_1) & \neg R \vee P \\ (C_2) & \neg Q \vee \neg P \\ (C_4) & R \vee P \end{array}$$

Exercise 5. In what follows we consider the theory of arrays defined in the lecture. We assume that the theory of indices \mathcal{T}_i is $LI(\mathbb{Z})$, and the theory of elements \mathcal{T}_e is $LI(\mathbb{Q})$.

Which of the formulae below are in the array property fragment and which are not? Justify your answer. (The universally quantified variables i, j are of sort index; the indices $k, l_i, u_i, i = 1, 2$ which are not universally quantified are considered to be constants of sort index)

- (1) $\forall i (a[a[i]] > a[i])$
- (2) $\forall i (i > a[i])$
- (3) $\forall i (a[i] > b[i])$
- (4) $\forall i (i \leq a[k] \rightarrow a[i] = a[k])$
- (5) $\forall i, j (l_1 < i < u_1 < l_2 < j < u_2 \rightarrow a[i] \leq a[j])$
- (6) $\forall i, j (l_1 < i < j < u_2 \rightarrow a[i] \leq a[j])$
- (7) $\forall i, j (l_1 < i \leq j < u_2 \rightarrow a[i] \leq a[j])$

Exercise 6.

(**Note:** the probability that such an exercise would come up in the exam is extremely low)

Consider the array property formula:

$$F : write(a, l, v_1)[k] = b[k] \wedge b[k] = v_2 \wedge a[k] = v_1 \wedge v_1 \neq v_2 \wedge \forall i (i \leq l-1 \rightarrow a[i] = b[i]) \wedge \forall i (i \geq l+1 \rightarrow a[i] = b[i])$$

- (1) Apply Steps 1–6 described in the lecture to F . Let F_6 be the formula obtained after Step 6.
- (2) Check the satisfiability of F_6 using one of the versions of the $DPLL(\mathcal{T})$ procedure presented in the class. For theory reasoning in combinations of theories use the Nelson-Oppen procedure.